

BASICS OF CLIMATE CHANGE


DR. RENEE A. MCPHERSON

ASSOCIATE PROFESSOR, GEOGRAPHY & ENVIRONMENTAL SUSTAINABILITY
UNIVERSITY DIRECTOR, SOUTH CENTRAL CLIMATE SCIENCE CENTER
UNIVERSITY OF OKLAHOMA



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

THE CLIMATE UMBRELLA



**Climate
Variability**

**Climate
“Normals”**

**Climate
Change**

**Climate
Extremes**



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

WEATHER VS. CLIMATE



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER




WEATHER VS. CLIMATE

Weather – **state of the atmosphere** with respect to heat or cold, wetness or dryness, calm or storm, clearness or cloudiness; **short-term**

Climate – **statistical collection of weather conditions** at a place over a period of years; **long-term**

“Climate is what you expect. Weather is what you get.”

CLIMATE VARIABILITY VS. CHANGE




© Jones & Bartlett Learning




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

REALLY LONG-TERM CLIMATE CHANGE


Shape of Earth's orbit
changes over
~100,000 yrs



Wobble (precession)
of Earth on its axis
changes over
~22,000 yrs



Tilt of Earth's
axis changes
over ~41,000 yrs



© Jones & Bartlett Learning





SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

NATURAL CAUSES OF CLIMATE CHANGE

External factors: (1) **luminosity** of sun (amount of incoming radiation), (2) Earth's **orbital mechanics** (tilt, precession, orbit shape) & (3) comet, meteorite, or asteroid **impact event**


Internal factors: (1) **plate tectonics** (location of land, weathering), (2) **ocean** temperatures & currents, & (3) natural changes in **atmospheric composition**



TEMPERATURE VS. LUMINOSITY OF SUN

Sun's energy output has been decreasing over past few decades

Energy from the sun cycles with sunspot activity (~11 yrs)



KEY POINTS

No single weather event (e.g., early autumn blizzard, December heat wave, landfalling hurricane) **is a sign of climate change**, but a **higher frequency** of certain events or trend toward **higher intensity events may be**.

There are natural drivers to climate change, but they typically **occur over 1000s to 100,000s of years**. Faster changes, like those that occur with volcanic eruptions, usually last only a few years and are part of climate variability, not climate change.




FROM THE 4TH NATIONAL CLIMATE ASSESSMENT (2017)

“Since NCA3 [Third National Climate Assessment], **stronger evidence has emerged for continuing, rapid, human-caused warming of the global atmosphere and ocean.** This report concludes that ‘it is **extremely likely** that **human influence has been the dominant cause** of the observed warming since the mid-20th century. For the warming over the last century, **there is no convincing alternative explanation** supported by the extent of the observational evidence.’” – Climate Science Special Report



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

4TH NATIONAL CLIMATE ASSESSMENT (2018)



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

WHY ARE IPCC REPORTS IMPORTANT?

IPCC = Intergovernmental Panel on Climate Change

Rigorous & transparent review process focusing on climate change, its impacts, and our ability to adapt and mitigate

Main assessment reports **summarize almost 10,000 peer-reviewed scientific papers** in both an easy-to-read format (for policymakers) & in a detailed manner (for researchers)

Policy relevant but not policy prescriptive



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

REASONS FOR CONCERN

Aggregate impacts & damages

**Risks of large-scale
discontinuities &
disruptions**



**Uneven distribution
of climate change
impacts**

**Risks of extreme
weather events**

**Risks to unique &
threatened systems**




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

IPCC: GLOBAL WARMING OF 1.5°C (OCT. 2018)

Five Reasons For Concern (RFCs) illustrate the impacts and risks of different levels of global warming for people, economies and ecosystems across sectors and regions.

Impacts and risks associated with the Reasons for Concern (RFCs)



IPCC 2018



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

KEY POINT

The **Intergovernmental Panel on Climate Change (IPCC)** reports and the **National Climate Assessment (NCA)** are the **two best sources** for expert assessments of climate change worldwide and across the United States.

Available at:

<http://www.ipcc.ch> (IPCC)

<http://www.globalchange.gov> (NCA)



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

The Greenhouse Effect

Some solar radiation is reflected by the Earth and the atmosphere.


Some of the infrared radiation passes through the atmosphere. Some is absorbed and re-emitted in all directions by greenhouse gas molecules. The effect of this is to warm the Earth's surface and the lower atmosphere.

Most radiation is absorbed by the Earth's surface and warms it.

Atmosphere


Earth's surface

Infrared radiation is emitted by the Earth's surface.




Natural Greenhouse Effect

Human Enhanced Greenhouse Effect




WE'VE KNOWN THIS FACT FOR 160 YEARS



“The atmosphere admits of the entrance of the solar heat, but checks its exit; and the **result is a tendency to accumulate heat at the surface of the planet.**” — John Tyndall, 1859

“**Doubling of CO₂ would raise surface temperature** by 5-6°C, or 9-11°F, above pre-industrial temperatures.” — Svante Arrhenius, 1896



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

KEY POINT


Greenhouse gases are necessary for Earth to be livable, **but adding too much into the atmosphere will disrupt** the long-term energy balance, **increasing the thermal energy** in the atmosphere.





CLIMATE CHANGE NOT FROM OZONE HOLE

Greenhouse gases selectively absorb infrared energy & convert it to heat energy to warm troposphere (surface to 12 miles)

Ozone selectively absorbs ultraviolet energy & converts it to heat energy to warm stratosphere (13 to 30 miles above surface)



ANTARCTIC OZONE HOLE (2018 & PAST)




Data from NASA's Aura satellite



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

IMPACT OF MONTREAL PROTOCOL



NASA says ozone hole stabilizing but won't fully recover until 2070


December 12, 2013 | By Tony Barboza

Email Share +1 38 Tweet 0

Los Angeles Times

The hole in the ozone layer is stabilizing but will take until about 2070 to fully recover, according to new research by NASA scientists.

The assessment comes more than two decades after the [Montreal Protocol](#), the international treaty that banned chlorofluorocarbons and other compounds that deplete the ozone layer, which shields the planet from harmful ultraviolet rays.



The ozone hole, shown in October, has stopped growing since the mid-1990s... (Ozone Hole Watch)



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER



KEY POINTS

The **ozone holes** at the North and South Poles are **NOT related to climate change!!**

Because of international cooperation, the **ozone layer is healing** and projected to **recover by 2070**




GLOBAL CARBON DIOXIDE BUDGET



GREENHOUSE GASES INCREASING

Significant increases in carbon dioxide, methane, & nitrous oxide observed since the industrial revolution

Changes in Greenhouse Gases from Ice-Core and Modern Data




© 2012 Pearson Education, Inc.




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

OBSERVATIONS OF CARBON DIOXIDE (CO₂)


Charles Keeling first measured CO₂ at the Mauna Loa Observatory, leading the scientific community to notice the human contribution to the greenhouse effect



GREENHOUSE GAS EMISSIONS BY GAS



Largest contributor is CO₂ from fossil fuel use




KEY POINT


Greenhouse gases have been increasing at an abnormally fast rate, primarily as a result of human activity (i.e., fossil fuel use, deforestation, agriculture, etc.). Our oceans and ecosystems cannot absorb CO₂ quickly enough to keep up. Except for water vapor, **these greenhouse gases are long-lived and will remain in our atmosphere for decades.**



Indicators of a Warming World




TECHNIQUES TO OBSERVE THE CLIMATE



Creative Commons



USGS



Kathy Krucker



Nat'l Park Service







Carlye Calvin



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

OBSERVATIONAL EVIDENCE FOR A WARMING CLIMATE




IPCC AR5



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

Land-Surface Temperature Anomaly

1°C




Data source: Berkeley Earth daily TAVG full dataset (experimental)
Global land-surface temperature anomaly
Base period: 1880-1920
<https://berkeleyearth.org/>

Antti Lipponen (@anttilip)
CC BY 4.0

SURFACE AIR TEMPERATURE OBSERVATIONS


Globally, the past three decades have been successively warmer, on average, than the prior decades. Multi-decadal warming is superimposed on decadal and interannual variability from natural climate patterns.




CLIMATE CHANGE IMPACTS ARE REGIONAL

For example, surface temperature changes are not uniform

Surface Temperature Change




Change in Temperature (°F)




NCA4, Vol 1


Annual Temperature



Winter Temperature



Summer Temperature





NCA4, Vol 1



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

REGIONAL CHANGES HAVE CAUSES




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER


CLIMATE CHANGE IMPACTS ARE REGIONAL

Precipitation & sea-level rise changes are not uniform



Annual Precipitation



Winter Precipitation



Spring Precipitation




NCA4, Vol 1



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

RAPID DECLINE OF ARCTIC SEA ICE



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

MOUNTAIN GLACIAL RETREAT




Okpilak Glacier 1907

Okpilak Glacier 2004




**National Snow and
Ice Data Center**





SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

AS OCEANS ABSORB CO₂, THEY BECOME MORE ACIDIC



NCA 2014; modified from Feely et al. 2009



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

WARMING OCEANS BLEACH CORALS



NOAA


Coral reefs endangered by bleaching in global event, researchers say

By Adam Dunnakey, CNN

Updated 1:06 PM ET, Thu October 8, 2015




NOAA CRW 5-km Night-Only HotSpot Year-to-date Maximum 08 Oct 2016




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

WARMER WATER + MELTING GLACIERS = SEA-LEVEL RISE



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

MANY OTHER CLIMATE CHANGE-RELATED CHANGES



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER


KEY POINTS

Historical observations demonstrate **rapid (decadal) climate changes** in surface temperature, sea ice, mountain glaciers, sea level, and other parts of our climate system.

These changes **are all consistent with a warming planet** resulting from increased greenhouse gases.

The changes **are not consistent with long-term natural variations** in our climate.

GLOBAL CLIMATE MODEL (GCM)



Models based on physical laws and statistical representations of observations


Provides reasonable description of physical changes, not detailed predictions

© 2012 Pearson Education, Inc.




SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

REAL VS. COMPUTED CLIMATE



Multi-Satellite Image Animation




Global Climate Model Simulation



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

USING GLOBAL CLIMATE MODELS

Using computer models from 20 climate modeling groups worldwide, global climate models that include both natural forcing and forcing from human activities best relate to the actual observations



IPCC AR5



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

FUTURE SCENARIOS IN IPCC & NCA


Several scenarios are used to depict how greenhouse gas (GHG) emissions may change in the future.

RCP 8.5 – GHG emissions continue to increase (“business as usual”)

RCP 6 – GHG emissions stabilized around 2100 using various technologies & reduction strategies


RCP 4.5 – similar to RCP 6 but with a lower stabilization target

RCP 2.6 – a “peak-and-decline” scenario where GHG emissions are reduced significantly over time




PROJECTED CHANGES IN AVERAGE ANNUAL TEMPERATURE


Lower Scenario (RCP4.5)



Higher Scenario (RCP8.5)




Change in Temperature (°F)



Projections for the
late 21st century
relative to 1976–2005



PROJECTED CHANGES IN SEASONAL PRECIPITATION



NCA4, Vol 1



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER

KEY POINTS


General trends over time and large-scale regional patterns **can be projected** using numerical climate models.

These **climate models** have **satisfactorily depicted changes** in the past.

Model results depicting future **changes in temperatures** **are more certain** than results depicting future changes in precipitation.

“DOWNSCALING” THE GCM OUTPUT

What is downscaling? – a method to use “low resolution” global climate model output (e.g., 100-500 km grid) & obtain “high resolution” (e.g., 10-50 km grid) climate projections



WHY DOWNSCALE?


Climate scientists downscale GCM data **to help answer stakeholders' questions** about how the climate will change in their location (i.e., impact assessments) & better represent local climates



Mountain climates




Coastal climates




Urban-rural climates



ENSEMBLE APPROACH TO DOWNSCALING



ENSEMBLE APPROACH TO DOWNSCALING



QUESTIONS?



SOUTH CENTRAL
CLIMATE ADAPTATION SCIENCE CENTER