



**SOUTH CENTRAL**  
CLIMATE ADAPTATION SCIENCE CENTER

# **Drought History**

For Louisiana's 9 Regions

**Updated 2021**

# In This Report

---

<b>What is drought?</b>	<b>3</b>
<b>Why be prepared for drought?</b>	<b>3</b>
<b>How to prepare for drought?</b>	<b>4</b>
<b>How is drought measured?</b>	<b>4</b>
<b>Climate Division 1: Northwest</b>	<b>7</b>
<b>Climate Division 2: North Central</b>	<b>10</b>
<b>Climate Division 3: Northeast</b>	<b>13</b>
<b>Climate Division 4: West Central</b>	<b>16</b>
<b>Climate Division 5: Central</b>	<b>19</b>
<b>Climate Division 6: East Central</b>	<b>22</b>
<b>Climate Division 7: Southwest</b>	<b>25</b>
<b>Climate Division 8: South Central</b>	<b>28</b>
<b>Climate Division 9: Southeast</b>	<b>31</b>
<b>Our Changing Climate</b>	<b>34</b>
<b>Sources of Drought Information &amp; Tools</b>	<b>37</b>

## What is drought?

Defining drought can be difficult because the impacts associated with drought are often far-reaching and devastating. A *meteorological drought* is a prolonged period when precipitation is below “normal” for the location (Heim 2002). An *agricultural drought* occurs when soils are too dry to grow healthy vegetation, particularly crops or forests. As water becomes scarce in rivers, lakes, and other water bodies, a *hydrological drought* develops. If, at any time, the water demands of society (e.g., water for drinking, maintaining lawns and gardens, washing clothes) exceed the availability of good-quality water, then a *socioeconomic drought* has occurred. A socioeconomic drought may arise even during times of normal precipitation because of increased water demand from a growing population, increased temperatures and wind speeds, new businesses, or other societal changes.

## Why be prepared for drought?

Since 1980, the National Oceanic and Atmospheric Administration has identified 29 droughts nationwide as weather disasters based on both damages and costs in the amount of \$269.6 billion dollars; 12 of which directly affected Louisiana. The 2012 drought, which at its height affected over 80% of the contiguous U.S., resulted in estimated damages and costs of over \$50 billion from both direct and indirect impacts.

Drought can result in crop, pasture, and forest damage; increased livestock and wildlife mortality; increased Fire hazard; threats to aquatic and wildlife habitats; increased water demand; and reduced water supplies.

Proper management of water resources is necessary to protect supplies for drinking water, sanitation, and Fire protection as well as to maintain economic activity and environmental sustainability. ***Because disasters affect families, neighbors, and businesses locally, community-level planning is necessary to reduce the vulnerability to drought in Louisiana.***

## ***“Droughts-of-Record” in Louisiana***

For purpose of planning, we consider the “drought-of-record” to be the drought with the worst environmental conditions rather than the drought with the worst recorded impacts. Hence, a shorter and less severe drought with high monetary losses in our recent past (e.g., during 2011) will not outweigh a long and severe drought in our early history, when fewer people lived in the region. We choose to prepare for the worst.

## How to prepare for drought?

Local officials and other key stakeholders in Louisiana will be better prepared for drought when they complete the following: (1) have assessed their vulnerability to drought, (2) understand past droughts and the local climate, (3) monitor drought, (4) prepare a thorough set of actions to be taken before, during, and after a drought, and (5) educate citizens on this plan.

Having a plan in place will enable these individuals to understand key factors to monitor so they may respond proactively to drought conditions early. Following this plan helps reduce the risk such that, when drought conditions occur, water resources do not run out. This report will help governmental officials and resource managers in Louisiana by overviewing the climate and drought history since record-keeping began in the late 19th century.

## How is drought measured?

To quantify drought severity, the scientific community has developed several methods to assess drought, including departure from normal precipitation, the Palmer Drought Severity Index, and the Standardized Precipitation Index. All three use weather observations to diagnose drought conditions. The simplest of these is the annual departure from normal precipitation, which is the actual precipitation total for the year subtracted from the annual normal. Large negative values indicate a precipitation deficit for that year.

*The Palmer Drought Severity Index* uses observations or estimates of precipitation, temperature, and soil water content. Values typically range from +4 representing extremely wet conditions to -4 representing extremely dry conditions. Values less than -1 indicate some level of drought, and the values become more negative with less rainfall and hotter temperatures.

The Palmer Drought Severity Index helps to diagnose agricultural drought because it is sensitive to soil moisture conditions and works well at relatively long-time scales. The index does not account for reservoir levels and streamflow, so it has drawbacks for diagnosing hydrological drought.

*The Standardized Precipitation Index* is based solely on precipitation but has the advantage of multiple time scales (e.g., 3 months, 6 months, 1 year) to better highlight short-term versus long-term droughts. Values typically range from +2 as extremely wet to -2 as extremely dry, with values less than -1 representing drought.


A more recent method to measure drought intensity is the U.S. Drought Monitor (Figure 1). This product depicts weekly drought conditions for the United States on a drought intensity scale of D0 to D4, with D0 representing areas that are abnormally dry and D4 representing areas of exceptional drought. Although the levels are subjectively determined, they are established through expert review of weather and water data, including local observations (e.g., NM Climate Center), as well as reports of drought impacts from local, tribal, state, and federal officials as well as the public and media. Figure 1 displays the weekly percentage of area in each of Louisiana's regions by D0 through D4 drought since 2000<sup>1</sup>.

# U.S. Drought Monitor

July 20, 2021

(Released Thursday, Jul. 22, 2021)

Valid 8 a.m. EDT




**Figure 1.** Example map of the U.S. Drought Monitor from the drought assessment issued for the week preceding June 10, 2021. The color scale (yellow to dark red) displays the level of drought from D0 (abnormally dry) to D4 (exceptional drought). Significant regional impacts on agriculture are designated with an “S” and regional impacts on water supply are designated with an “L”. The maps are released each Thursday at 8:30 a.m. Eastern Time. Courtesy of the National Drought Mitigation Center.

## ***Has Louisiana experienced drought?***

***Drought is a recurring condition in Louisiana*** and is part of our climate. Our climate history can provide us insight into what we may see in the future. Being “drought ready” means, in part, that we recognize how our climate has changed over time. This report examines Louisiana’s drought History.

### **The Climate of Louisiana**

Temperature and precipitation are the two main elements of our climate. Because Louisiana is located in the middle latitudes, and north of the Gulf of Mexico, its citizens experience a wide range of weather conditions. Hence, our climate highly variable, from year to year, season to season, and month to month.



### **Louisiana Climate Divisions**


- 1. Northwest**
- 2. North Central**
- 3. Northeast**
- 4. West Central**
- 5. Central**
- 6. East Central**
- 7. Southwest**
- 8. South Central**
- 9. Southeast**

***Figure 2. Climate Divisions of Louisiana***




## Climate Division 1: Northwest

Northwest Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 3 shows the annual temperature (top) and annual precipitation (bottom) in northwest Louisiana since 1895. The annual temperature for northwest Louisiana averages 64.7 degrees Fahrenheit, while precipitation averages 49.96 inches. Warmer-than-average periods have spanned the 1920s through the mid-1950s and the late 1990s through the early 2010s. Significant periods of drier-than-average conditions include the mid-1890s through the mid-1900s, the 1950s, the 1960s through the early 1970s, and the early 2010s.




**Figure 3.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in northwest Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for Northwest Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 4. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 4).



**Figure 4. Top graph:** The monthly average temperature (in degrees Fahrenheit) across Northwest Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across Northwest Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.





**Figure 5.** Above is the Palmer Drought Severity Index for Northwest Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 2: North Central

North central Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 6 shows the annual temperature (top) and annual precipitation (bottom) in north-central Louisiana since 1895. The annual temperature for north central Louisiana averages 64.22 degrees Fahrenheit, while precipitation averages 53.86 inches. Warmer-than-average periods have spanned the 1920s through the 1950s. Significant periods of drier-than-average conditions include the mid-1890s through the mid-1900s, the early 1950s, the 1960s through the early 1970s, and the early 2010s.




**Figure 6.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in north-central Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for North Central Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 7. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 7).



**Figure 7. Top graph:** The monthly average temperature (in degrees Fahrenheit) across North Central Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across North Central Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.


### Palmer Drought Severity Index for North Central Louisiana



**Figure 8.** Above is the Palmer Drought Severity Index for North Central Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 3: Northeast

Northeast Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 9 shows the annual temperature (top) and annual precipitation (bottom) in northeast Louisiana since 1895. The annual temperature for northeast Louisiana averages 64.69 degrees Fahrenheit, while precipitation averages 54.03 inches. Warmer-than-average periods have spanned the 1920s, the mid-1940s through the 1950s, and the late 1990s through the early 2010s. Significant periods of drier-than-average conditions include the mid-1890s through the mid-1900s, the 1950s through the early 1970s, and the early 2010s.




**Figure 9.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in northeast Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for Northeast Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 10. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 10).



**Figure 10. Top graph:** The monthly average temperature (in degrees Fahrenheit) across Northeast Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across Northeast Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.





**Figure 11.** Above is the Palmer Drought Severity Index for Northeast Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 4: West Central

West central Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 12 shows the annual temperature (top) and annual precipitation (bottom) in west-central Louisiana since 1895. The annual temperature for west central Louisiana averages 65.62 degrees Fahrenheit, while precipitation averages 53.23 inches. Warmer-than-average periods have spanned the 1920s, the mid-1930s through the late 1930s, the mid-1940s through the mid-1950s, and the late 1990s through the early 2010s. Significant periods of drier-than-average conditions include the mid-1890s through the mid-1900s, the mid-1910s, the mid-1950s through the early 1970s, and the early 2010s.




**Figure 12.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in west-central Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for West Central Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 13. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 13).



**Figure 13. Top graph:** The monthly average temperature (in degrees Fahrenheit) across West Central Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across West Central Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.



**Figure 14.** Above is the Palmer Drought Severity Index for West Central Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 5: Central

Central Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 15 shows the annual temperature (top) and annual precipitation (bottom) in central Louisiana since 1895. The annual temperature for central Louisiana averages 66.46 degrees Fahrenheit, while precipitation averages 57.31 inches. Warmer-than-average periods have spanned the 1920s through the mid-1950s and the late 1990s through the early 2010s. Significant periods of drier-than-average conditions include the mid-1890s through the mid-1900s, the mid-1910s, the mid 1950s through the early 1970s, the late 1990s to the early 2000s, and the early 2010s




**Figure 15.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in central Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for Central Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 16. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 16).



**Figure 16. Top graph:** The monthly average temperature (in degrees Fahrenheit) across Central Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across Central Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.






**Figure 17.** Above is the Palmer Drought Severity Index for Central Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 6: East Central

East central Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 18 shows the annual temperature (top) and annual precipitation (bottom) in east-central Louisiana since 1895. The annual temperature for east central Louisiana averages 66.80 degrees Fahrenheit, while precipitation averages 61.65 inches. Warmer-than-average periods have spanned the 1920s, the late 1940s through the 1950s, and the mid-1990s through the early 2010s. Significant periods of drier-than-average conditions include the late 1890s through the mid-1900s, the mid to late 1910s, the mid to late 1930s, the early 1960s through the early 1970s, and the late 1990s through the early 2010s




**Figure 18.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in east-central Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for East Central Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 19. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 19).



**Figure 19. Top graph:** The monthly average temperature (in degrees Fahrenheit) across East Central Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across East Central Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.



**Figure 20.** Above is the Palmer Drought Severity Index for East Central Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than  $-4$  being categorized as extreme drought. Graph readings between  $-2$  and  $-3$  indicate moderate drought.


## Climate Division 7: Southwest

Southwest Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 21 shows the annual temperature (top) and annual precipitation (bottom) in southwest Louisiana since 1895. The annual temperature for southwest Louisiana averages 67.98 degrees Fahrenheit, while precipitation averages 58.21 inches. Warmer-than-average periods have spanned the late 1900s, the 1920s through the 1930s, the mid-1940s through the 1950s, the mid-1990s through the early 2010s. Significant periods of drier-than-average conditions include the 1900s, the mid to late 1910s, the 1930s, the late 1940s through the mid-1950s, the early 1960s through the early 1970s, and the late 2000s through the early 2010s.




**Figure 21.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in southwest Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for Southwest Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 22. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 22).



**Figure 22. Top graph:** The monthly average temperature (in degrees Fahrenheit) across Southwest Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across Southwest Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.





**Figure 23.** Above is the Palmer Drought Severity Index for Southwest Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 8: South Central

South central Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 24 shows the annual temperature (top) and annual precipitation (bottom) in south-central Louisiana since 1895. The annual temperature for south central Louisiana averages 68.10 degrees Fahrenheit, while precipitation averages 59.81 inches. Warmer-than-average periods have spanned the 1920s through the 1930s, the mid-1940s through the 1950s, and the mid-1990s through the early 2010s. Significant periods of drier-than-average conditions include the 1900s and the 1910s, the late 1930s, the 1950s through the early 1970s, and the late 1990s through the early 2010s.




**Figure 24.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in south-central Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.

To understand when there is the greatest stress on water availability for South Central Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 25. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 25).



**Figure 25. *Top graph:*** The monthly average temperature (in degrees Fahrenheit) across South Central Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. ***Bottom graph:*** The average total precipitation by month across South Central Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.



**Figure 26.** Above is the Palmer Drought Severity Index for South Central Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.


## Climate Division 9: Southeast

Southeast Louisiana has experienced a wide range of temperatures and precipitation over the past several decades. Abnormally hot and dry conditions have occurred multiple times since the early 1900s. Figure 27 shows the annual temperature (top) and annual precipitation (bottom) in southeast Louisiana since 1895. The annual temperature for southeast Louisiana averages 69.0 degrees Fahrenheit, while precipitation averages 61.03 inches. Warmer-than-average periods have spanned the 1920s through the 1930s, the mid-1940s through the 1950s, the mid-1990s through the early 2010s. Significant periods of drier-than-average conditions include the late 1890s through the late 1910s, the 1930s, the 1950s, the mid-1960s, and the late 1990s through the early 2010s.



**Figure 27.** The average annual temperature (top graph) and total annual precipitation (bottom graph) in southeast Louisiana from 1895 to 2021. To highlight warmer, cooler, wetter, or drier periods, 5-year moving averages are shaded. On the top graph, red shading (above the horizontal line) indicates warmer periods and blue shading (below the line) notes cooler periods than average. Similarly, on the bottom graph, green shading (above the horizontal line) highlights wetter periods and brown shading (below the line) highlights drier periods than average. Extended periods of relatively warm temperatures or low precipitation are outlined in red boxes.


To understand when there is the greatest stress on water availability for Southeast Louisiana, the average monthly temperature and precipitation, as well as their average highest and lowest monthly values, are shown in Figure 28. Warmer temperatures result in greater water loss by evaporation and transpiration. The warmest temperatures typically occur during July and August (top of Figure 28.)



**Figure 28. Top graph:** The monthly average temperature (in degrees Fahrenheit) across Southeast Louisiana using data from 1895 to 2021. The green line is the average of all climate-division average temperatures for that time period. The red line is the highest monthly average, and the blue line is the lowest. **Bottom graph:** The average total precipitation by month across Southeast Louisiana using data from 1895 to 2021. The blue bar is the highest monthly precipitation; the green is the average precipitation total recorded for that month; the gold is the lowest precipitation total recorded for that month.



### Palmer Drought Severity Index for Southeast Louisiana



**Figure 29.** Above is the Palmer Drought Severity Index for Southeast Louisiana. Note how any regions below the zero line indicate dry conditions, with any instance less than -4 being categorized as extreme drought. Graph readings between -2 and -3 indicate moderate drought.

## Our Changing Climate

As documented by the Intergovernmental Panel on Climate Change (IPCC), the Earth's climate has warmed during the past 100 years. Climatic changes are expected to impact water supplies. Although the annual totals of rainfall may remain fairly constant, or even slightly increase, there may be less water available for public consumption. This is possible if precipitation is more intense and less frequent, water may need to be discharged from reservoirs to avoid flooding. Intense runoff during extreme precipitation events will likely pollute the water, rendering large volumes unsuitable for human consumption.


To investigate what the future might look like in Louisiana, we use global climate models (GCMs) to project future conditions based on trends of global emissions. The global climate models use physics to calculate temperature, moisture, wind, pressure, etc. at each point in a grid that covers the globe. GCMs are not used to predict the weather on any given day; rather, they help us understand how weather on average might change. Typically, a single grid cell in a GCM covers an area of 100-500 km (62-310 miles). This can make it difficult to obtain useful information at regional-to-local scales. Using downscaling techniques, we can take the coarse resolution (i.e., 100-500 km area) of the GCM to a finer resolution that provides detailed information at more regional-to-local scales. We've used statistical downscaling techniques, rather than dynamical downscaling techniques, to obtain finer resolution climate projections for our region (i.e., 10 km (6 miles) area).

Statistical downscaling refers to a group of downscaling techniques that build a relationship between the regional/local (represented by historical observations) and global scales using statistical methods. Most statistical downscaling approaches begin by comparing GCM output for a particular time period in the past with historical observations during that same historical time period. By comparing GCM output and actual climate observations, our researchers can see the relationships between global and regional climate patterns, and ultimately describe these relationships using statistics. Historical observations often include temperature and precipitation measurements from networks like the Oklahoma Mesonet, the Community Collaborative Rain, Hail, & Snow Network (CoCoRaHS), NOAA's Cooperative Observer Network, etc.


Emissions scenarios are used to show how future trajectories of key greenhouse gases may impact the climate system. For our datasets, we show how the climate responds to the low scenario (e.g., a drastic reduction in global emissions) and the high scenario (e.g., business as usual).

Figures 30 and 31 present mid-century (2036-2065) and end-of-century (2070-2099) downscaled climate projections for drought related events across the south-central U.S. Each figure includes projections for a low emissions scenario on the left and a high emissions scenario on the right.

## Mid-Century Projected Change in the Average Annual Longest Dry Spell Length




## End-of-Century Projected Change in the Average Annual Longest Dry Spell Length




**Figure 30:** The length of the longest dry spell in Louisiana is expected to be six days for the mid-century low emissions scenario. This affected region is small and concentrated in the northwest corner of the state. However, in the mid-century high emissions scenario, this dry spell duration extends along the whole state's western border. As one can see, the magnitude of these annual average dry spells is significantly greater in the high emissions scenario compared to the low emissions scenario, regardless of which part of the century's projections you are examining.

## Mid-Century Projected Change in the Average Annual Number of Days with Rain



## End-of-Century Projected Change in the Average Annual Number of Days with Rain



**Figure 31:** These projections show how the yearly average number of days with rain will be affected over time. The mid-century low emissions scenario projection shows miniscule change overall, while its corresponding high emissions scenario projection predicts approximately a 4 day decrease in number of days with rain for much of the state. These reductions in rainfall are exacerbated in the end of the century high emissions scenario, with some parts of Louisiana seeing as much as a ten day decrease in the number of days with rain.

# Drought Resources

There are several excellent sources of information for drought monitoring and reporting, ranging from national to local (Table 1). These resources are designed to help you plan and prepare for drought conditions in your region.

Table 1 summarizes several excellent sources of information for drought monitoring and reporting, ranging from national to local.

Table 1: Sources of Drought Information & Tools

Source	Web Address	Uses
<b>Major Sources of Information</b>		
National Integrated Drought Information System	<a href="http://drought.gov/">drought.gov/</a>	Consolidated source of drought information, monitoring & reporting tools, including many of the other sources listed below
National Drought Mitigation Center	<a href="http://drought.unl.edu/">drought.unl.edu/</a>	Consolidated source of drought information, including drought planning, monitoring reporting, risks, and impacts
State Climate Office - Louisiana	<a href="http://losc.lsu.edu">losc.lsu.edu</a>	Consolidated source of Louisiana climate information
<b>Specific Drought-Related Tools</b>		
U.S. Drought Monitor: <i>National Drought Mitigation Center</i>	<a href="http://droughtmonitor.unl.edu">droughtmonitor.unl.edu</a>	Current and past diagnoses of drought conditions, both nearby and across the United States
U.S. Seasonal Drought & Precipitation Outlooks: <i>Climate Prediction Center</i>	<a href="http://cpc.ncep.noaa.gov/products/expert_assessment/sdo_summary.php">cpc.ncep.noaa.gov/products/expert_assessment/sdo_summary.php</a>	Large-scale trends in drought across the U.S. for the next few months; Expert assessments (not forecasts) of possible changes in precipitation conditions over a range of times (6-10 days, 8-14 days, 1 month, & 3 months)

## Bibliography

Heim, R. R., 2002: A Review of Twentieth-Century Drought Indices Used in the United States. *Bulletin of the American Meteorological Society*, **83**, 1149-1165.

NOAA National Centers for Environmental Information (NCEI) U.S. Billion-Dollar Weather and Climate Disasters (2017).  
<https://www.ncdc.noaa.gov/billions/>